

Volumes by Shells


Thomas R. Cameron

September 5, 2025

1 Shell Method

Sometimes the method of disks and washers is not the most efficient way to find the volume of a solid of revolution.

For example, consider the solid of revolution formed by taking the plane region bounded by $y = \sqrt{x}$, $x = 1$, $x = 4$, and the x -axis, and revolving that region around the y -axis. Note that each cross section is a washer; however, the inner radius of the washer is $x = 1$, for $0 \leq y \leq 1$, and is $x = y^2$, for $1 \leq y \leq 2$. Hence, to find the volume of the solid of revolution using the washer method we need two integrals.



The shell method uses cylindrical shells to approximate the volume of a solid of revolution. A cylindrical shell is a solid enclosed by two circular cylinders. If the outer cylinder has radius r_2 and the inner cylinder has radius r_1 , then the volume of the shell is given by

$$\begin{aligned} V &= \pi (r_2^2 - r_1^2) h \\ &= \pi (r_2 + r_1)(r_2 - r_1) h \\ &= 2\pi \left(\frac{r_1 + r_2}{2} \right) (r_2 - r_1) h. \end{aligned}$$

Now, divide the interval $[a, b]$ into n subintervals of length Δx . Then, the plane region has been divided into n strips, which we denote by R_1, R_2, \dots, R_n . When revolved around the y -axis, these strips generate tube-like solids S_1, S_2, \dots, S_n that are nested one inside the other, see Figure 1.




Figure 1: Shell Method, split the region into n strips and revolve around y -axis

Note that these strips have curved upper bounds. However, if the strips are thin we can approximate their area using a rectangle. These rectangles, when revolved around the y -axis, will produce cylindrical shells whose volume closely approximate the volume of the tubes, see Figure 2.

To implement the shell method, consider the subinterval $[x_{k-1}, x_k]$ and let x_k^* denote the midpoint of this interval. If we construct a rectangle of height $f(x_k^*)$ over this interval, then revolving the rectangle around




Figure 2: Shell Method, approximate the area of each strip by a rectangle and revolve around y -axis

the y -axis will produce a cylindrical shell of average radius x_k^* , height $f(x_k^*)$, and thickness Δx . Hence, the volume of this shell is given by

$$V_k = 2\pi x_k^* f(x_k^*) \Delta x.$$

Therefore, we can approximate the volume of the solid of revolution as follows

$$V \approx 2\pi \sum_{i=1}^n x_i^* f(x_i^*) \Delta x.$$

Taking the limit as $n \rightarrow \infty$ gives us

$$V = 2\pi \lim_{n \rightarrow \infty} \sum_{i=1}^n x_i^* f(x_i^*) \Delta x = 2\pi \int_a^b x f(x) dx.$$